Mouse mAb to human CDNF (clone 6G5)

Catalogue #	302-100
Immunogen:	Human CDNF
Immunogen Description:	Recombinant human CDNF protein produced using CHO-based Icosagen Cell factory Ltd. proprietary suspension cell line. Purified from cell culture supernatant
Alternative Names:	ARMETL1, cerebral dopamine neurotrophic factor, ARMET-like protein 1, conserved dopamine neurotrophic factor
Uniprot ID:	Q49AH0
Clonality:	Mouse monoclonal
Clone:	6G5
Class:	mIgG1
Reactivity:	Human, no reactivity with mouse CDNF Binds to the C-terminal part of the human CDNF (aa 126-187)
Application:	ELISA, WB, IF, IHC
Protocol:	Monoclonal antibody working titer has to be established practically for each particular antigen and assay format
Buffer:	PBS pH 7.4, with 0.1% sodium azide
Shipping:	Shipping in ambient conditions.
Storage:	Store at -15…-25 °C. Avoid multiple freeze-thaw cycles.
Background:	CDNF is a trophic factor for midbrain dopamine neurons in vivo. It prevents the 6-OHDA- (Lindholm et al. 20007; Voutilainen et al., 2011) and MPTP-induced degeneration (Airavaara et al., 2012) of dopamine neurons in rodent models of Parkinson’s disease. When administered after 6-OHDA or MPTP –lesioning it restores the dopaminergic function and prevents degeneration

of dopamine neurons in substantia nigra pars compacta



Figure 1. Western Blot testing of anti-CDNF monoclonal antibody (6G5). Line 1. PageRuler Prestained Protein Ladder (#SM0671 Fermentas). Line 2. Recombinant CDNF expressed into the supernatant of CHO cell culture medium.



Figure 2. Immunohistochemistry testing of anti-CDNF monoclonal antibody 6G5. Analysis was performed using formalin-fixed paraffin-embedded human cerebral cortex tissue sections from Alzheimer's disease patients. Tissue sections were boiled with sodium citrate buffer (pH 6) for antigen retrieval. Incubation with primary antibody at 5 µg/ml was performed overnight at 4°C. DAKO EnVisionTM Detection System, Peroxidase/DAB was used for visualization. Sections were counterstained with toluidine blue and mounted with Eukitt mounting medium. A. CDNF staining by monoclonal antibody 6G5; B. Negative staining without primary antibody.



Figure 3. Immunofluorescence detection of human CDNF expressed in U2OS cells. CDNF was visualized using anti-CDNF antibody clone 6G5 at 1 µg/ml. Goat anti-mouse AlexaFluor488 was used as secondary antibody. For nuclear staining DAPI was used. ArrayScan VTI platform (Thermo Scientific) was used for image acquisition (10x objective). Composite picture was generated using pseudocolors green for CDNF specific signal and blue for nuclei. A. CDNF-expressing U2OS cells; B. Negative control (non-transfected U2OS cells).

(for reference only; data shown are from a single production batch at release and may not be representative of other batches)